Volume 32 Number 2

ANTELOPE SPECIALIST GROUP

September 2015

In this Issue...

Gnusletter Editor

Å This issue: S. Shurter

New Research antE ¹ ‘ ‘ ‘

New Research and Reports

Trade of ungulates in Moroccan markets for decoration



Table of cities in which ungulate parts were observed showing coordinates of the city, number of repeat surveys in the markets for 2013/3014 and numbers of each species observed 2013/2014

Bergin D. and Nijman V. (2014a) Ongoing, open wildlife trade in Moroccan markets: reptiles, mammals and their derivatives. *TRAFFIC Bulletin* xx: xx-xx

Bergin D. and Nijman V. (2014b) Illegal and open wildlife trade in Moroccoôs capital. *SWARA* July ï September

Cuzin, F., Sehhar, E. A. and Wacher, T. (2007) Etude pour lô®laboration de lignes directrices et dôun plan dôaction strat®gique pour la conservation des ongul®s au Maroc. Haut Commissariat aux Eaux et For°ts et ¨ la Lutte Contre le D®sertiýcation (HEFLCD), Projet de Gestion des Aires Prot®g®es (PGAP) et Banque Mondiale, Global Environ ĭ ʻ 6 M Gȯ uɐpouɹ ʻ V

Be ĭ ʻ " ! M Gȯ ʻ ʻ Eaux et For°ts et ¨ la

cus, Hylochoerus meinertzhageni). The red ï þanked duiker *C. rufýlatus* is represented in the PNDD by one of its southernmost area of occurrence in Cameroon (Fig. 2). The buffalo (*Syncerus caffer*) has not been observed during the surveys but informal information suggests that it might occur in the parkôs northern sector. Tracks and

The Red-fronted gazelle (Eudorcas ruĀfrons) in the Boundou Community Nature Reserve, Senegal.

Cl®ment[1] C., Ceulers[1] V., Medina[2] R., Welle[2] M., Rabeil[3]* T. & Jullien[4] E.
**Corresponding author : Thomas Rabeil, thomas.rabeil@sahara-conservation.org*
[1] International Independent Consultant
[2] CORENA - Association of the Boundou Community Nature Reserve, Tambacounda, BP 238, SENEGAL
[3] Sahara Conservation Fund
[4] Joseph Fourier Grenoble University

The Red-fronted gazelle (*Eudorcas ruýfrons*) is a Sahelian migratory species which occurs in grasslands and open savannas, from Senegal to Sudan. This gazelle is listed as Vulnerable in the IUCN Red List, threatened by habitat degradation, competition with domestic livestock and poaching (UICN, 2008).

However, we are concerned that its conservation assessment as Vulnerable may be too optimistic. A continuing decline in their populations has been reported since the last assessment in 2008 and this trend has been conýrmed recently in Niger and Chad (Ahmat Brahim Siam, Director of the Protected Areas in Chad, Hamissou Halilou Malam Garba, Head of Protected areas unit in Niger; personal communication, 2015). As a result, the Convention on Migratory Species has recently decided to include the red-fronted gazelle on CMS Appendix I (IUCN, 2014). Despite the rising concern of the scientiýc community for this species, there are still very few data and studies to help reassessing the red-fronted gazelle status and taking appropriate conservation measures.

In Senegal, the red-fronted gazelle is present in the Boundou Community Nature Reserve, a 1,200 kmĮ protected area located in Eastern Senegal, between the Ferlo Reserve and Niokolo Koba National Park. In this Sahelo-Sudanian environment, the other ungulate species observed are mainly the common warthog (*Phacochoerus africanus*), common duiker (*Sylvicapra grimmia*), oribi (*Ourebia ourebi*) and roan antelope (*Hippotragus equinus*). The most frequent carnivores are spotted hyena (*Crocuta crocuta*) and common jackal (*Canis aureus*). Some felids are also present but rare, such as serval (*Leptailurus serval*), caracal (*Caracal caracal)*, lion (*Panthera leo*) and leopard (*Panthera pardus*).

The Boundou Reserve, ofýcially recognized as a Community Nature Reserve in 2009 by conversion of a tourist hunting area, is managed by local communities through an association (CORENA) with the support of the General Council of Isere (France) and the regional technical services. A technical team hired by the CORENA and a local eco-guards network are working together, implementing the area surveillance, wildlife monitoring and conservation actions. For several years, the red-fronted gazelle has beneýted from closer attention to its status as it has been identiýed as an umbrella species for this reserve. A ýrst survey of *Eudorcas ruýfrons* based on foot transects was carried out in 2011, and this initiative developed subsequently into more general large mammal monitoring.

Currently, the survey is based on 12 foot transects of 5 km spread across the reserve. The transects were set up according to the different types of vegetation and the main water sources of the reserve in order to get a representative sample of the main habitats. This protocol is implemented with limited resources but is easily replicated 3 times a year (hot, rainy and cool seasons). The foot survey is conducted by local rangers, known as eco-guards, and technical advisers from CORENA. In 2014, the survey was carried out twice, once during the hot season (May) and once in the rainy season (July). Two camera-traps were also used in an opportunistic way during the dry season, but unfortunately one of them was destroyed by a spotted hyena. Finally, during the transects, fecal samples of red-fronted gazelle were collected and sent to the WildGenes Laboratory of the Royal Zoological Society of Scotland, for genetic analysis in order to assess the taxonomy of the different red-fronted gazelle subspecies.

The preliminary results from last yearôs transect survey provided positive news about the local gazelle population. Results in 2014 indicate that the red-fronted gazelle is well distributed all over the reserve during the hot and the rainy seasons. The contact rate per



Localization of the 12 foot transects in the Boundou Community Nature Reserve, S®n®gal.



A red-fronted gazelle (male) taken by a camera trap in the Boundou Reserve, Senegal

IUCN, 2008. IUCN Red List. Available from https://www.iuc-nredlist.org (accessed May 2014)
IUCN, 2014MMMMMMMM ŝ ᴰ ᵐ Mʻ ˢ ή Mʻ ¯ᵛ ž 1 î MQuito, Ecuador, Novem]œ 4-9, 2014M

[6]Udzungwa Ecological Monitoring Centre, c/o Udzungwa Mountains National Park, PO Box 99, Mangôula, Tanzania
[7]Department for Food and Resource Economy, Faculty of Science,

Resource use and the nutritional status of sable antelope in the Okavango Delta region of northern Botswana

Michael C. Hensman[1]

Unsustainable hunting threatens both biodiversity and local livelihoods. Despite high levels of research effort focused on understanding the dynamics of bushmeat trade and consumption, current research islargely site speciýc. Without synthesis and quantitative analysis of available case studies, the national and regional characteristics of bushmeat trade and consumption remain largely speculative, impeding efforts to inform national and regional policy on bushmeat trade. Here we describe the structure and content of the West and Central African bushmeat database which holds quantitative data on bushmeat sales, consumption and offtake for 177 species from 275 sites across 11 countries in two regions, spanning three decades of research. Despite this wealth of available data, we found important biases in research effort. The majority of studies in West and Central Africa have collected market data, which although providing a useful record of bushmeat sales, are limited in their ability to track changes in hunting offtake. In addition, few data exist for West Africa, and few studies have tracked changes over time, using repeat sampling. With new initiatives in the regions to track bushmeat hunting, this database represents an opportunity to synthesise current and future data on bushmeat hunting, consumption and trade in West and Central Africa, identify gaps in current understanding, and systematically target future monitoring efforts.

Conservation, hunting, exploitation, wild meat

** Corresponding authors at: School of Life Sciences, University of Sussex, Brighton BN1 9QG, UK. Tel.: +44 (0)1273 873502 (J.P.W. Scharlemann). Environmental Change Institute, School of Geography, University of Oxford, Oxford OX1 3QY, UK. Tel.: +44 (0)1223 363835 (L.M Coad). E-mail addresses: gemma.taylor.14@ucl.ac.uk (G. Taylor), j.scharlemann@sussex.ac.uk (J.P.W. Scharlemann), Marcus.Rowcliffe@ioz.ac.uk (M. Rowcliffe), no ξ i, ẋ ʾ es (Q erstanding, and systematically tg n*

Cephalophus zebra / Zebra Duiker

One complete skin in good condition collected by B ¦ TZLER at Sound®dou on 30/01/95. The animal was caught in a steel snare to the north of Sound®dou, near the [Ziam ᵐ ɤ ƀ p M ᴺ ɓ ɤ ᵐ Î ɹ ᴀ l M ᵀ ᵠ Į Ů p ᵀ א ᵐ M

VOL. 32 NO. 2

Dorcas Gazelles on Buri Peninsula *photo by David Mallon*

Steve Shurter, Editor
White Oak Conservation Holdings
581705 White Oak Road
Yulee FL 32097

www.whiteoakwildlife.org